Buthod, A.K. and B.W. Hoagland. 2013. New to Oklahoma: Murdannia keisak (Commelinaceae). Phytoneuron 2013-93: 1–3. Published 3 December 2013. ISSN 2153 733X

NEW TO OKLAHOMA: MURDANNIA KEISAK (COMMELINACEAE)

AMY K. BUTHOD

Oklahoma Biological Survey/ Oklahoma Natural Heritage Inventory/Robert Bebb Herbarium University of Oklahoma 111 East Chesapeake Norman, Oklahoma 73019 amybuthod@ou.edu

BRUCE W. HOAGLAND Oklahoma Natural Heritage Inventory/ Department of Geography and Environmental Sustainability University of Oklahoma Norman, Oklahoma 73019

ABSTRACT

This paper documents the occurrence of a vascular plant taxon previously unknown to the flora of Oklahoma. Murdannia keisak, an Asian member of the Commelinaceae, was discovered in the Mountain Fork River in southeastern McCurtain County.

KEY WORDS: Murdannia, Commelinaceae, Oklahoma, McCurtain County, exotic

Murdannia keisak (Hassk.) Hand.-Mazz. (Commelinaceae; watermoving herb) is a weedy annual forb native to eastern Asia (Faden 2000; Seward 1958). In North America it was first reported in 1935 from South Carolina (Hotchkiss 1940, 1951), although the New York Botanic Garden herbarium has a Louisiana collection from 1927 (Dunn & Sharitz 1990a). It has since been found throughout the southeastern USA, as well as in Oregon and in Washington, where it is listed as a noxious weed (BONAP 2013; USDA, NRCS 2013; Washington Administrative Code 2005). Murdannia keisak has also been found in Europe (Faden 2000). The collections reported here are the first for the state of Oklahoma (Hoagland et al. 2004).

Vouchers. USA. Oklahoma. McCurtain Co.: On the Lower Mountain Fork River, Presbyterian Falls area, T5S R26E Sec. 31, 29 Aug 2013, Buthod & Hoagland AB-10591 (OKL); on the Lower Mountain Fork River, Presbyterian Falls area, T5S R26E Sec. 31, 4 Oct 2013, Buthod & Hoagland AB-10592 (OKL).

In the USA, Murdannia keisak is found in wet areas including river and creek margins, tidal marshes, swamps, and ditches (Faden 2000; Hotchkiss 1951; Rundell & Diamond 1999). It is thought to have been a contaminant of rice and is frequently found in old rice fields (Dunn & Sharitz 1990a).

A sterile specimen of Murdannia keisak (AB-10591) was first collected in August 2013 on the lower Mountain Fork River in southeastern McCurtain County. Associated species included Gratiola brevifolia, Hydrocotyle verticillata, Taxodium distichum, Xyris jupicai, and the endangered Harperella nodosa. A second, fruiting collection (AB-10592) was made approximately 1.0 km upstream in October 2013. Associated species included Impatiens capensis, Itea virginica, Polygonum persicaria, Sacciolepis striata, and Taxodium disticum. Plants of Murdannia at both sites numbered in the hundreds (Fig. 1).




Figure 1. Population of *Murdannia keisak* on the Mountain Fork River, McCurtain County, Oklahoma. Photo by Amy Buthod

Although rice was once grown in McCurtain County (Reasoner 1974), the presence of *Murdannia keisak* in Oklahoma may be attributed to waterfowl. *M. keisak* can produce 9,000-70,000 seeds/m², and they have been found in great abundance in the stomachs of ducks (Dunn and Sharitz 1990a; Hotchkiss 1940, 1951). It is also possible that pieces of the plant floated in from elsewhere; adventitious roots are produced by larger plants at the nodes, allowing for vegetative reproduction via fragmentation (Dunn and Sharitz 1990b; Ferrero et al. 2012).

According to Newberry (1991), Murdannia keisak may reduce rates of water flow because of its rhizomatous growth and fibrous roots. It grows fast and forms a thick mat, allowing it to outcompete native vegetation (Ferrero et al. 2012). It has also been shown to easily adapt to different environmental conditions (Dunn & Sharitz 1991). The Oklahoma population of Murdannia keisak will need to be carefully monitored. The Forest Service sensitive species Calamovilfa arcuata K.E. Rogers and Vernonia lettermanii Engelm. ex A. Gray are found on the Mountain Fork River, as is the western-most known population of the endangered Harperella nodosa (Rose) Rose (Buthod & Hoagland 2013; Hoagland et al. 2004; U.S. Forest Service 2005). We intend to evaluate the full extent of M. keisak in the Mountain Fork drainage in conjunction with future work on the endangered Harperella nodosa.

LITERATURE CITED

Buthod, A.K. and B.W. Hoagland. 2013. Noteworthy collections: Oklahoma. Castanea 78: 213-215.

- Dunn, C.P. and R.R. Sharitz. 1990a. The history of *Murdannia keisak* (Commelinaceae) in the southeastern United States. Castanea 55: 122–129.
- Dunn, C.P. and R.R. Sharitz. 1990b. The relationship of light and plant geometry to self-thinning of an aquatic annual herb, *Murdannia keisak* (Commelinaceae). New Phytol. 115: 559-565.
- Dunn, C.P. and R.R. Sharitz. 1991. Population structure, biomass allocation, and phenotypic plasticity in *Murdannia keisak* (Commelinaceae). Amer. J. Bot. 78: 1712–1723.
- Faden, R.B. 2000. Commelinaceae. Pp. 170–197, in Flora of North America Editorial Committee (eds.). Flora of North America North of Mexico, Vol. 22. New York and Oxford.
- Ferrero, A.F. Tesio, M. Tabacchi and F. Vidotto. 2012. The effects of water management, timing and the rate of several herbicides on the growth of *Murdannia keisak* (Hassk.) Handel-Mazz. Crop Protect. 38: 53–56.
- Hoagland B.W., A.K. Buthod, I.H. Butler, P.H.C. Crawford, A.H. Udasi, W.J. Elisens, and R.J. Tyrl. 2004. Oklahoma Vascular Plants Database. http://www.oklahomaplantdatabase.org/
- Hotchkiss, N. 1940. Range extensions of marsh and aquatic plants. Rhodora 42: 20–22.
- Hotchkiss, N. 1951. Range extensions of marsh and aquatic plants. 2. Rhodora 53: 91–93.
- BONAP. 2013. The Biota of North America Program. ">http://www.bonap.net/>
- Newberry, G. 1991. Factors affecting the survival of the rare plant, *Sagittaria fasciculata* E.O. Beal (Alismataceae). Castanea 56: 59-64.
- Reasoner, R.C. 1974. Soil survey, McCurtain County, Oklahoma. USDA Soil Conservation Service, Washington, D.C.
- Rundell, H. and A. Diamond. 1999. Noteworthy collections from Alabama. Castanea 64: 355–356. Seward, A.N. 1958. Manual of Vascular Plants of the Lower Yangtze Valley, China. Oregon State College, Corvallis.
- USDA, NRCS. 2013. The PLANTS Database. National Plant Data Team, Greensboro, North Carolina. http://plants.usda.gov> Accessed January 2013.
- U.S. Forest Service. 2005. Forest Service Sensitive Species that are not listed or proposed under the

ESA. <http://www.fs.fed.us/biology/resources/pubs/tes/fs_ss_1dec04.pdf> Washington Administrative Code. 2005. State noxious weed list and schedule of monetary penalties, chapter 16-750. State of Washington, Olympia.